Novel Coronavirus (COVID-19)

Recommendations for Use of Personal Protective Equipment

Routes of Transmission

Ontario Medical Association

- Cases and clusters demonstrate that COVID-19 is transmitted through the Droplet/Contact route
- Most cases are linked to person-to-person transmission through close contact
- There is no evidence that COVID-19 is transmitted through the airborne route except possibly in special circumstances

Key Recommendations for Care of Patients with Suspected or Confirmed COVID-19

When providing routine care:	 Use Droplet and Contact precautions Isolate the patient in a single room 	 *Aerosol-generating medical procedures Endotracheal intubation, including during
When planning or anticipating to conduct aerosol- generating medical procedures*:	 Use Airborne precautions Place the patient in an airborne infection isolation room (AIIR) If AIIR is not available, isolate the patient in a single room with a closed door 	 cardio-pulmonary resuscitation Cardio-pulmonary resuscitation Open airway suctioning Bronchoscopy (Diagnostic or Therapeutic) Surgery and autopsy Sputum induction (Diagnostic or Therapeutic) Non-invasive positive pressure ventilation for acute respiratory failure (CPAP, BiPAP3- 5) High flow oxygen therapy Note: The collection of a nasopharyngeal swab is <u>not</u> an aerosol generating procedure.

Precautions

Risk Assessments

- Health Care Workers have a responsibility to perform a Point-of-Care risk assessment before every patient interaction
- Organizations have a responsibility to conduct an Organizational Risk Assessment and education/train staff accordingly
- Organizations should apply a <u>hierarchy of hazard controls</u>, which include elimination and substitution, engineering and systems controls, administrative controls, and personal protective equipment.

Droplet and Contact Precautions

- Gloves
- Protective eyewear
- Surgical/procedure mask
- Isolation gown

Airborne Precautions

- N95 Respirator
- AIIR

Healthcare	Providers and Support Staff	Contact Precautions	Precautions	required	otier
In-Patient Healthcar	e Facilities				
Patient's room	Providing direct care to suspect or confirmed COVID-19 patients (including nasopharyngeal and oropharyngeal swab collection)	~			
Patient's room	Aerosol-generating procedures performed on suspect or confirmed COVID-19 patients	~	~		
Patient transit areas	Any activity that does not involve contact with suspect or confirmed COVID-19 patients				Routine practices and additional precautions based on risk assessment
Triage	Preliminary screening not involving direct contact	~			PPE not required if separated by physical barrier or able to maintain 2m distance from patient
Administrative areas	Administrative tasks that do not involve contact with patients			\checkmark	
Ambulatory and Out	patient Healthcare Facilities				
Consultation room	Physician examination of patient with suspected or confirmed COVID-19	~			
Administrative areas	Administrative tasks that do not involve contact with patients			~	
Reception/Triage	Preliminary screening not involving direct contact	~			PPE not required if separated by physical barrier or able to maintain 2m distance from patient
Home Care					
In patient's home	Visiting clients/patients suspected or confirmed to have COVID-19	\checkmark			
Long Term Care Hom	ne				
Any location	Providing direct care to suspect or confirmed COVID-19 residents (including nasopharyngeal and oropharyngeal swab collection)	~			
Any location	Providing CPAP and/or open suctioning to suspect or confirmed COVID-19 residents		~		
Administrative areas	Administrative tasks that do not involve contact with resident suspected or confirmed to have COVID-19			~	

Droplet &

Airborne

No PPE

Other

Visitors and Environmental Service Workers

All Care Settings			
Patient's room	Entering the room of a suspect or confirmed COVID-19 person	~	Visitors should be discouraged or limited
	(including entering to clean after and between consultations)		

Patients Suspected or Confirmed to have COVID-19

Triage	Provide patient with surgical/procedure mask if tolerated.
	Ask patient to maintain 2m distance or separation by a physician barrier.
	Ask patient to perform hand hygiene.
Ambulatory and (Dutpatient Healthcare Facilities
Triage/Reception	Provide patient with surgical/procedure mask if tolerated.
	Ask patient to maintain 2m distance or separation by a physician barrier.
Waiting room	Provide patient with surgical/procedure mask if tolerated.
	• Immediately move the patient to a single patient room or separate area away from others. If this is not feasible, ensure special distance of at least 2m from patient
Consultation	Provide patient with surgical/procedure mask if tolerated.
room	Ask patient to perform hand hygiene.